

Zagreb Polynomials, Co-Polynomials of Bipartite and Strongly Bipartite Graphs

Chandra Prakash Singh

Ex. Head, Department of Physics, Sunderrao Solanke, Mahavidyalaya Majalgaon Dist. Beed (M.S.) India
 rautnk87@gmail.com

Abstract: First Zagreb polynomial of a graph G with vertex set $V(G)$ and edge set $E(G)$ is defined as $M_1(G,x) = \sum_{uv \in E(G)} x^{d_u + d_v}$ and the first Zagreb index can be obtained from it as $M_1(G) = \frac{\partial M_1(G,x)}{\partial x} |_{x=1}$. In this paper Zagreb polynomials, co-polynomials and corresponding Zagreb indices, co-indices are obtained for path, cycle, double and strong double graphs.

Keywords: Cycle graph, double graph, path graph, strong double graph, Zagreb co-polynomial, Zagreb index, Zagreb polynomial

1. Introduction

Let G be a simple, finite, connected graph with vertex set $V(G)$ and edge set $E(G)$. The degree of a vertex $u \in V(G)$ is denoted by d_u and is the number of vertices adjacent to u . The edge connecting the vertices u and v is denoted by uv . A molecular graph is presentation of the structural formula of a chemical compound in terms of graph theory whose vertices correspond to the atoms of compound and edges correspond to chemical bonds [1]. The topological index is a numerical parameter mathematically derived from the graph structure, several such topological indices have been considered in theoretical chemistry and have found some applications in QSPR/QSAR study [2].

A complement \bar{G} of a graph G consist of the same set of vertices, where two vertices v and w are adjacent by an edge vw if and only if they are not adjacent in G . Hence $vw \in E(\bar{G}) \Leftrightarrow vw \notin E(G)$. A complement graph consists of a number of edges and the degree of vertex v which are represented as $\bar{m} = \binom{n}{2} - m$ and $d_{\bar{G}}(v) = n-1-R_G(v)$ respectively. If the graph G has n vertices and m edges, then double graph $D[G]$ has $2n$ vertices $4m$ edges. The first, second Zagreb indices and geometric-arithmetic index of double and strong double graph of path and cycle graphs were studied in [3]. The double graph of G is constructed by considering two copies of G in which a vertex v_i in first copy is adjacent to a vertex v_j in the second copy of v_i and v_j [4]. The strong double is a double in which a vertex v_i in the copy is adjacent to a vertex v_j in the second copy if $i = j$. It is denoted as $SD[G]$ [5]. Zagreb indices and multiplicative Zagreb indices of double graphs of subdivision graphs were investigated by M. Togan et al. [6].

Degree based topological indices of strong double graphs have been studied in [7-8]. The relations between some Zagreb indices and Zagreb co-indices of graphs can be seen in papers [9-10]. General fifth Zagreb polynomial of benzene ring were studied in [11]. Li and W. Gao expressed the fourth Zagreb polynomial and sixth polynomial in terms eccentricity of a graph [12]. The Zagreb group indices and polynomials are studied in [13]. Zagreb indices and Zagreb polynomials have been studied in [14]. Distance based topological indices of double and strong double graphs were

computed by Mirajkar [15]. The properties of strong double graphs of spanning trees of G were found in [16].

The Zagreb polynomials are defined as [17-18]

$$\begin{aligned} M_1(G,x) &= \sum_{uv \in E(G)} x^{d_u + d_v}, & (1) \\ M_2(G,x) &= \sum_{uv \in E(G)} x^{d_u \times d_v}, & (2) \\ M_3(G,x) &= \sum_{uv \in E(G)} x^{|d_u - d_v|}, & (3) \\ M_4(G,x) &= \sum_{uv \in E(G)} x^{d_u(d_u + d_v)}, & (4) \\ M_5(G,x) &= \sum_{uv \in E(G)} x^{d_v(d_u + d_v)}, & (5) \\ \bar{M}_1(G,x) &= \sum_{uv \notin E(G)} x^{(d_u + d_v)}, & (6) \\ \bar{M}_2(G,x) &= \sum_{uv \notin E(G)} x^{(d_u \times d_v)}. & (7) \\ \bar{M}_3(G,x) &= \sum_{uv \notin E(G)} x^{|d_u - d_v|}, & (8) \\ \bar{M}_4(G,x) &= \sum_{uv \notin E(G)} x^{d_u(d_u + d_v)}, & (9) \end{aligned}$$

And

$$\bar{M}_5(G,x) = \sum_{uv \notin E(G)} x^{d_v(d_u + d_v)} \quad (10)$$

The first Zagreb index can be calculated as the first derivative of first Zagreb polynomial, $x = 1$ by [19]

$$M_1(G) = \frac{\partial M_1(G,x)}{\partial x} |_{x=1}. \quad (11)$$

All the symbols and notations used in this paper are standard and taken mainly from books of graph theory [20-22]. In this paper we study Zagreb polynomials, co-polynomials of double, strong double graphs of path graph P_4 and cycle graph C_4 .

2. Materials and Methods

Let G be a finite, simple, connected graph with vertex set $V(G)$ and edge set $E(G)$. Molecular graph is a graph in which the vertices correspond to the atoms and the edges to the bonds of a molecule. Molecular graphs of path graph P_4 , double graph $D[P_4]$, strong double graph $SD[P_4]$ and cycle graph C_4 , double graph of C_4 and strong double graph of C_4 are shown figure (1-2). Degree d_u of a vertex u is the number of vertices adjacent to u . The order and size of strong double $SD[G]$ are $2n$ and $4m+n$ respectively. The degree of a vertex v in $SD[G]$ is $\text{deg}_{SD[G]}(v) = 2\text{deg}_G(v) + 1$. If the graph has n vertices and the m edges, then the double graph $D[G]$ has $2n$ vertices and $4m$ edges, in particular $\text{deg}_{D[G]}(v,2) = 2\text{deg}_G(v)$. The complement of G denoted by \bar{G} , is a simple graph on the same set of vertices $V(G)$ in which two vertices

u and v are adjacent, that is connected by an edge uv , iff they are not adjacent in G . Hence $uv \in \bar{G} \Leftrightarrow uv \notin E(G)$. Degree of a vertex u in G is denoted d_u ; the degree of the same vertex in \bar{G} is given by $deg_{\bar{G}}(u) = n-1-d_u$. Degrees of vertices of path, cycle, double, strong double graphs are used in finding the Zagreb polynomials and co-polynomials.

3. Results and Discussion

It is observed from molecular graph of path graph P_4 , there are $|E_{1,2}| = n-2$ and $|E_{2,3}| = n-3$ edges. For double graph of P_4 the edge partition is $|E_{2,4}| = n$ and $|E_{4,4}| = n-4$ and in strong double graph of P_4 , it is $|E_{3,4}| = n$, $|E_{4,4}| = n-4$ and $|E_{3,3}| = n-6$. The Zagreb polynomials, co-polynomials of P_4 , $D[P_4]$ and $SD[P_4]$ are computed as follows.

Path graph P_4

Theorem 1.1. First Zagreb polynomial of P_4 is $(n-2)x^3 + (n-3)x^4$.

Proof. First Zagreb polynomial of P_4

$$\begin{aligned} M_1(G, x) &= \sum_{u \in V(G)} x^{d_u + d_v} \\ &= |E_{1,2}|x^{1+2} + |E_{2,2}|x^{2+2} \\ &= (n-2)x^3 + (n-3)x^4. \end{aligned}$$

Theorem 1.2. Second Zagreb polynomial of P_4 is $(n-2)x^2 + (n-3)x^4$.

Proof. Second Zagreb polynomial of P_4

$$\begin{aligned} M_2(G, x) &= \sum_{u \in V(G)} x^{d_u \times d_v} \\ &= |E_{1,2}|x^{1 \times 2} + |E_{2,2}|x^{2 \times 2} \\ &= (n-2)x^2 + (n-3)x^4. \end{aligned}$$

Theorem 1.3. Third Zagreb polynomial of P_4 is $(n-2)x$

Theorem 1.4. Fourth Zagreb polynomial of P_4 is $(n-2)x^3 + (n-3)x^8$.

Theorem 1.5. Fifth Zagreb polynomial of P_4 is $(n-2)x^6 + (n-3)x^8$.

Theorem 1.6. First Zagreb co-polynomial of P_4 is $(n-2)x^3 + (n-3)x^2$.

Proof. First Zagreb co-polynomial of P_4

$$\begin{aligned} \bar{M}_1(G, x) &= \sum_{u \in \bar{E}(G)} x^{d_u + d_v} \\ &= |E_{1,2}|x^{2+1} + |E_{2,2}|x^{1+1} \\ &= (n-2)x^3 + (n-3)x^2. \end{aligned}$$

Theorem 1.7. Second Zagreb co-polynomial of P_4 is $(n-2)x^2 + (n-3)x$.

Proof. Second Zagreb co-polynomial of P_4

$$\begin{aligned} \bar{M}_2(G, x) &= \sum_{u \in \bar{E}(G)} x^{d_u \times d_v} \\ &= |E_{1,2}|x^{2 \times 1} + |E_{2,2}|x^{1 \times 1} \\ &= (n-2)x^2 + (n-3)x. \end{aligned}$$

Theorem 1.8. Third Zagreb co-polynomial of P_4 is $(n-2)x + (n-3)$

Theorem 1.9. Fourth Zagreb co-polynomial of P_4 is $(n-2)x^6 + (n-3)x^2$.

Theorem 1.10. Fifth Zagreb co-polynomial of P_4 is $(n-2)x^3 + (n-3)x^2$.

Theorem 2.1. First Zagreb polynomial of double graph of P_4 is $4nx^{12} + 4(n-4)x^{16}$.

Theorem 2.2. Second Zagreb polynomial of double graph of P_4 is $4nx^{32} + 4(n-4)x^{64}$.

Theorem 2.3. Third Zagreb polynomial of double graph of P_4 is $4n x^4$.

Theorem 2.4. Fourth Zagreb polynomial of double graph of P_4 is $4nx^{48} + 4(n-4)x^{128}$.

Proof. Fourth Zagreb polynomial double graph of P_4

$$\begin{aligned} M_4(D[G], x) &= \sum_{uv \in E(G)} x^{d_u + d_v} \\ &= 4 \sum_{uv \in E(D[G])} x^{2d_u + 2d_v} \\ &= 4|E_{2,4}|x^{2d_u + 2d_v} + 4|E_{4,4}|x^{2d_u + 2d_v} \\ &= 4nx^{48} + 4(n-4)x^{128}. \end{aligned}$$

Theorem 2.5. Fifth Zagreb polynomial of double graph of P_4 is $4nx^{48} + 4(n-4)x^{128}$.

Proof. Fifth Zagreb polynomial of double graph of P_4

$$\begin{aligned} M_5(D[G], x) &= \sum_{uv \in E(G)} x^{d_u \times d_v} \\ &= 4 \sum_{uv \in E(D[G])} x^{2d_u \times 2d_v} \\ &= 4|E_{2,4}|x^{2d_u \times 2d_v} + 4|E_{4,4}|x^{2d_u \times 2d_v} \\ &= 4nx^{48} + 4(n-4)x^{128}. \end{aligned}$$

Theorem 2.6. First Zagreb co-polynomial of double graph of P_4 is $4nx^6 + 4(n-4)x^{-2}$.

Theorem 2.7. Second Zagreb co-polynomial of double graph of P_4 is $4nx^{-3} + 4(n-4)x$.

Theorem 2.8. Third Zagreb co-polynomial of double graph of P_4 is $4nx^4 - (n-4)$.

Theorem 2.9. Fourth Zagreb co-polynomial of double graph of P_4 is $(4n)x^{40} + 4(n-4)x^{18}$.

Proof. Fourth Zagreb co-polynomial of double graph of P_4

$$\begin{aligned} \bar{M}_4(G, x) &= \sum_{u \in \bar{E}(G)} x^{d_u + d_v} \\ &= 4 \sum_{u \in \bar{E}(G)} x^{2d_u + 2d_v} \\ &= 4|E_{2,4}|x^{5(5+3)} + 4|E_{4,4}|x^{3(3+3)} \\ &= (4n)x^{40} + 4(n-4)x^{18}. \end{aligned}$$

Theorem 2.10. Fifth Zagreb co-polynomial of double graph of P_4 is $4nx^{24} + 4(n-4)x^{18}$.

Proof. Fifth Zagreb co-polynomial of double graph of P_4

$$\begin{aligned} \bar{M}_5(G, x) &= \sum_{u \in \bar{E}(G)} x^{d_u \times d_v} \\ &= 4 \sum_{u \in \bar{E}(G)} x^{2d_u \times 2d_v} \\ &= 4|E_{2,4}|x^{5(5+3)} + 4|E_{4,4}|x^{3(3+3)} \\ &= 4nx^{24} + 4(n-4)x^{18}. \end{aligned}$$

Theorem 3.1. First Zagreb polynomial of strong double graph of P_4 is $nx^{16} + (n-4)x^{18} + (n-6)x^{14}$.

Proof. First Zagreb polynomial of strong double graph of P_4

$$\begin{aligned}
M_1(SD[G],x) &= \sum_{uv \in E(G)} x^{d_u+d_v} \\
&= |E_{3,4}|x^{[2d_u+1]+[2d_v+1]} + |E_{4,4}|x^{[2d_u+1]+[2d_v+1]} + |E_{3,3}| \\
&\quad x^{[2d_u+1]+[2d_v+1]} \\
&= nx^{16} + (n-4)x^{18} + (n-6)x^{14}.
\end{aligned}$$

Theorem 3.2. Second Zagreb polynomial of strong double graph of P_4 is $nx^{63} + (n-4)x^{81} + (n-6)x^{49}$.

Proof. Second Zagreb polynomial of strong double graph of P_4

$$\begin{aligned}
M_2(SD[G],x) &= \sum_{uv \in E(G)} x^{d_u \times d_v} \\
&= |E_{3,4}|x^{[2d_u+1] \times [2d_v+1]} + |E_{4,4}|x^{[2d_u+1] \times [2d_v+1]} + |E_{3,3}| \\
&\quad x^{[2d_u+1] \times [2d_v+1]} \\
&= nx^{63} + (n-4)x^{81} + (n-6)x^{49}.
\end{aligned}$$

Theorem 3.3. Third Zagreb polynomial of strong double graph of P_4 is $nx^2 + (2n-10)$.

Theorem 3.4. Fourth Zagreb polynomial of strong double graph of P_4 is $nx^{112} + (n-4)x^{162} + (n-6)x^{98}$.

Theorem 3.5. Fifth Zagreb polynomial of strong double graph of P_4 is $nx^{144} + (n-4)x^{162} + (n-6)x^{98}$.

Theorem 3.6. First Zagreb co-polynomial of strong double graph of P_4 is $(n-6)x^{18} + nx^{16} + (n-4)x^{14}$.

Theorem 3.7. Second Zagreb co-polynomial of strong double graph of P_4 is $(n-6)x^{81} + nx^{63} + (n-4)x^{49}$.

Theorem 3.8. Third Zagreb co-polynomial of strong double graph of P_4 is $nx^2 + 2n - 10$.

Theorem 3.9. Fourth Zagreb co-polynomial of strong double graph of P_4 is $(n-6)x^{162} + nx^{144} + (n-4)x^{98}$.

Proof. Fourth Zagreb co-polynomial of double graph of P_4

$$\begin{aligned}
\overline{M}_4(SD[G],x) &= \sum_{u \in E(G)} x^{d_u(d_u+d_v)} \\
&= \sum_{u \in E(G)} x^{(2d_u+1)((2d_u+1)+(2d_v+1))} \\
&= |E_{3,3}|x^{(2d_u+1)((2d_u+1)+(2d_v+1))} \\
&\quad + |E_{3,4}|x^{(2d_u+1)((2d_u+1)+(2d_v+1))} + |E_{4,4}| \\
&\quad x^{(2d_u+1)((2d_u+1)+(2d_v+1))} \\
&= (n-6)x^{162} + nx^{144} + (n-4)x^{98}.
\end{aligned}$$

Theorem 3.10. Fifth Zagreb co-polynomial of strong double graph of P_4 is $(n-6)x^{162} + nx^{112} + (n-4)x^{98}$.

Proof. Fifth Zagreb co-polynomial of double graph of P_4

$$\begin{aligned}
\overline{M}_5(SD[G],x) &= \sum_{u \in E(G)} x^{d_u(d_u+d_v)} \\
&= |E_{3,3}|x^{(2d_v+1)((2d_u+1)+(2d_v+1))} \\
&\quad + |E_{3,4}|x^{(2d_v+1)((2d_u+1)+(2d_v+1))} + |E_{4,4}| \\
&\quad x^{(2d_v+1)((2d_u+1)+(2d_v+1))} \\
&= (n-6)x^{162} + nx^{112} + (n-4)x^{98}.
\end{aligned}$$

Cycle graph C_4

It is observed from molecular graph of cycle graph C_4 there are $|E_{2,2}| = n$ edges. For double graph of $D[C_4]$, the edge partition is $|E_{4,4}| = 2n$ and in strong double graph $SD[C_4]$ it is

$|E_{55}| = 3n$. The Zagreb polynomials, co-polynomials of cycle graph C_4 , $D[C_4]$ and $SD[C_4]$ are computed as follows.

Theorem 4.1. First Zagreb polynomial of C_4 is nx^4 .

Theorem 4.2. Second Zagreb polynomial of C_4 is nx^4 .

Theorem 4.3. Third Zagreb polynomial of C_4 is n .

Theorem 4.4. Fourth Zagreb polynomial of C_4 is nx^8 .

Theorem 4.5. Fifth Zagreb polynomial of C_4 is nx^8 .

Theorem 4.6. First Zagreb co-polynomial of C_4 is nx^2 .

Theorem 4.7. Second Zagreb co-polynomial of C_4 is nx .

Theorem 4.8. Third Zagreb co-polynomial of C_4 is n .

Theorem 4.9. Fourth Zagreb co-polynomial of C_4 is nx^2 .

Theorem 4.10. Fifth Zagreb co-polynomial of C_4 is nx^2 .

Theorem 5.1. First Zagreb polynomial of double graph of C_4 is $8x^{16}$.

Proof. First Zagreb polynomial of double graph of C_4

$$\begin{aligned}
M_1(D[G],x) &= \sum_{uv \in E(G)} x^{d_u+d_v} \\
&= 4|E_{4,4}|x^{2d_u+2d_v} \\
&= 8nx^{16}.
\end{aligned}$$

Theorem 5.2. Second Zagreb polynomial of double graph of C_4 is $8nx^{64}$.

Proof. Second Zagreb polynomial of double graph of C_4

$$\begin{aligned}
M_2(D[G],x) &= \sum_{uv \in E(G)} x^{d_u \times d_v} \\
&= 4|E_{4,4}|x^{2d_u \times 2d_v} \\
&= 8nx^{64}.
\end{aligned}$$

Theorem 5.3. Third Zagreb polynomial of double graph of C_4 is $8n$.

Theorem 5.4. Fourth Zagreb polynomial of double graph of C_4 is $8nx^{128}$.

Theorem 5.5. Fifth Zagreb polynomial of double graph of C_4 is $8nx^{128}$.

Theorem 5.6. First Zagreb co-polynomial of double graph of C_4 is $8nx^{12}$.

Theorem 5.7. Second Zagreb co-polynomial of double graph of C_4 is $8nx^{36}$.

Theorem 5.8. Third Zagreb co-polynomial of double graph of C_4 is $8n$.

Theorem 5.9. Fourth Zagreb co-polynomial of double graph of C_4 is $8nx^{72}$.

Proof. Fourth Zagreb co-polynomial of double graph of C_4

$$\begin{aligned}
M_4(D[G],x) &= \sum_{uv \in E(G)} x^{d_u(d_u+d_v)} \\
&= 4|E_{4,4}|x^{2d_u(2d_u+2d_v)} \\
&= 8nx^{72}.
\end{aligned}$$

Theorem 5.10. Fifth Zagreb co-polynomial of double graph of C_4 is $8nx^{72}$.

Proof. Fifth Zagreb co-polynomial of double graph of C_4

$$\begin{aligned}
M_5(D[G],x) &= \sum_{uv \in E(G)} x^{d_u(d_u+d_v)} \\
&= 4|E_{4,4}|x^{2d_v(2d_u+2d_v)} \\
&= 8nx^{72}.
\end{aligned}$$

Theorem 6.1. First Zagreb polynomial of strong double graph of C_4 is $(2n+4)x^{22}$.

Proof. First Zagreb polynomial of strong double graph of C_4

$$\begin{aligned}
M_1(SD[G],x) &= \sum_{uv \in E(G)} x^{d_u+d_v} \\
&= |E_{5,5}|x^{(2d_u+1)+(2d_v+1)}
\end{aligned}$$

$$= (2n + 4)x^{22}.$$

Theorem 6.2. Second Zagreb polynomial of strong double graph of C_4 is $(2n+4)x^{121}$.

Proof. Second Zagreb polynomial of strong double graph of C_4

$$\begin{aligned} M_2(SD[G],x) &= \sum_{uv \in E(G)} x^{d_u \times d_v} \\ &= |E_{5,5}|x^{(2d_u+1) \times (2d_v+1)} \\ &= (2n + 4)x^{121}. \end{aligned}$$

Theorem 6.3. Third Zagreb polynomial of strong double graph of C_4 is $2n+4$.

Theorem 6.4. Fourth Zagreb polynomial of strong double graph of C_4 is $(2n+4)x^{242}$.

Theorem 6.5. Fifth Zagreb polynomial of strong double graph of C_4 is $(2n+4)x^{242}$.

Theorem 6.6. First Zagreb co-polynomial of strong double graph of C_4 is $(2n+4)x^{-8}$.

Theorem 6.7. Second Zagreb co-polynomial of strong double graph of C_4 is $(2n+4)x^{16}$.

Theorem 6.8. Third Zagreb co-polynomial of strong double graph of C_4 is $(2n+4)$.

Theorem 6.9. Fourth Zagreb co-polynomial of strong double graph of C_4 is $(2n+4) x^{32}$.

Proof. Fourth Zagreb co-polynomial of strong double graph of C_4

$$\begin{aligned} \overline{M}_4(SD[G],x) &= \sum_{uv \notin E(G)} x^{d_u(d_u+d_v)} \\ &= |E_{5,5}| \sum_{uv \in E(G)} x^{(2d_u+1)+[(2d_u+1)+(2d_v+1)]} \\ &= (2n+4) x^{32}. \end{aligned}$$

Theorem 6.10. Fifth Zagreb co-polynomial of strong double graph of C_4 is $(2n+4) x^{32}$.

Proof. Fifth Zagreb co-polynomial of strong double graph of C_4

$$\begin{aligned} \overline{M}_5(SD[G],x) &= \sum_{uv \notin E(G)} x^{d_u(d_u+d_v)} \\ &= |E_{5,5}| \sum_{uv \in E(G)} x^{(2d_v+1)+[(2d_u+1)+(2d_v+1)]} \\ &= (2n+4) x^{32}. \end{aligned}$$

Zagreb indices and co-indices

1) First Zagreb index of double graph of P_4

$$\begin{aligned} M_1D[G] &= \frac{\partial M_1(D[G],x)}{\partial x} \Big|_{x=1} \\ &= (n-2)3+(n-3)4 \\ &= 7n-18. \end{aligned}$$

2) Fourth Zagreb co-index of double graph of C_4

$$\begin{aligned} \overline{M}_4D[G] &= \frac{\partial \overline{M}_4(D[G],x)}{\partial x} \Big|_{x=1} \\ &= 4(n^2+2n)162 \\ &= 648(n^2+2n). \end{aligned}$$




Figure 1: The double graphs of P_4 and C_4 .




Figure 2: The strong double graphs of P_4 and C_4 .

Table 1: Zagreb indices, co-indices of path, cycle, double, strong double graphs of P_4 and C_4 .

Zagreb indices → Graph ↓	M_1	M_2	M_3	M_4	M_5	\overline{M}_1	\overline{M}_2	\overline{M}_3	\overline{M}_4	\overline{M}_5
P_4	7n-18	6n-16	n-2	11n-30	14n-36	5n-12	3n-7	2n-5	8n-18	5n-12
$D[G]P_4$	112n-256	384n-256	16n	320n-512	704n-2048	56n+32	-(12n+16)	17n-4	232n-288	156n-288
$SD[G]P_4$	48n-156	193n-618	4n-10	372n-1236	306n-1236	48n-164	193n-682	4n-10	306n-1364	372n-1364
C_4	4n	4n	0	8n	8n	2n	N	0	2n	2n
$D[G]C_4$	128n	512n	0	1024n	1024n	96n	288n	0	576n	576n
$SD[G]C_4$	22(2n+4)	121(2n+4)	0	242(2n+4)	242(2n+4)	-8(2n+4)	16(2n+4)	0	32(2n+4)	32(2n+4)

4. Conclusion

Zagreb polynomials, co-polynomials and corresponding Zagreb indices of path, cycle, double, strong double graphs for path graph P_4 and cycle graph C_4 are studied.

References

[1] M. R. R. Kanna, S. Roopa and L. Parashivamurthy, Topological indices of Vitamin D₃, International Journal of Engineering and Technology, 7(4) (2018) 6276-6284.
 [2] V. R. Kulli, On Bannatti-Sombor indices, SSRG International Journal of Applied Chemistry, 8(1) (2001) 21-25.
 [3] M. Imran, S. Akhter, Degree based topological indices of double graphs and strong double graphs, Discrete

Mathematics and Applications, 9(5) (2017) 1750066-15.
 [4] E. Munarini, C. Perelli, A. Scaliola and N. Z. Salvi, Double graphs, Discrete Mathematics, 302(2002) 242-254.
 [5] V. M. Diudea, J. Chem. Inf. Comput. Sci., 37(1997) 292-299.
 [6] M. Togan, A. Yurttas, A. S. Cervic and N. Congul, Zagreb indices and multiplicative Zagreb indices of double graphs of subdivision graphs, TWMS Journal of Applied Engineering Mathematics, 9(2) (2019) 404-412.
 [7] M. Rafiullah, H. Muhammad, A. Siddiqui, M. K. Siddiqui and M. Dhlamini, On degree-based topological indices for strong double graphs, Hindawi, Journal of Chemistry, Volume 2021, Article Id-4852459, 12 pages.

- [8] M. S. Sardar, M. A. Ali, F. Asfrac and M. Cancan, On topological indices of double and strong double graphs of silicon carbide $Si_2C_3-I[p, q]$, *Euroasian Chem. Commun.*, 5(2023) 37-49.
- [9] K. Kiruthika, Zagreb indices and Zagreb co-indices of some graph operations, *International Journal of Advanced Research in Engineering and Technology*, 7(3) (2016) 25-41.
- [10] I. Gutmn, B. Furtula, Z. K. Vukicevic and G. Popivoda, On Zagreb indices and co-indices, *MATCH Mathematical and Computer Chemistry*, 74(2015) 5-16.
- [11] P. Sarkar, A. Pal, General fifth M-polynomials of benzene ring implanted in the P-type surface in 2D network, *Biointerface Research in Applied Chemistry*, 10(6) (2020) 6881-6892.
- [12] L. Yan, W. Gao, Eccentric related indices of an infinite class of nanostar dendrimers (II), *Journal of Chemical and Pharmaceutical Research*, 8(5) (2016) 359-362.
- [13] N. K. Raut, The Zagreb group indices and polynomials, *International Journal of Modern Engineering Research*, 6(10) (2016) 84-87.
- [14] M. R. Farahani, Zagreb indices and Zagreb polynomials of polycyclic aromatic hydrocarbons, *Journal of Chemical Acta*, 2(2013) 70-72.
- [15] K. Mirajkar, S. Konnur, Distance-based topological indices of double graphs and strong double graphs, *Ratio Mathematics*, Volume 48, 2023, 1-20.
- [16] T. A. Chisti, H. A. Ganie and S. Pirzada, Properties of strong double graphs, *Journal of Discrete Mathematical Science and Cryptography*, 4(17) (2014) 311-319.
- [17] A. U. Rehaman, W. Khalid, Zagreb polynomials and redefined Zagreb indices of line graph of $HAC_5C_6C_7[p, q]$ nanotube, *Open Journal of Chemistry*, 1(1) (2018) 26-35.
- [18] B. Basavanagoud, P. Jakkannavar, On the Zagreb polynomials of transformation graphs, *International Journal of Scientific Research in Mathematical and Statistical Sciences*, 5(6) (2018) 328-335.
- [19] M. R. Farahani, Zagreb indices and Zagreb polynomials of polycyclic aromatic hydrocarbons PAHs, *Journal of Chemical Acta*, 2(2013) 70-72.
- [20] Narsing Deo, *Graph Theory*, Prentice-Hall of India, New Delhi (2007).
- [21] N. Trinajstić, *Chemical Graph Theory*, CRC Press, Boca Raton, FL., 1992.
- [22] R. Todeschini, and V. Consonni, *Handbook of Molecular Descriptors*, Wiley-VCH:Weinheim, 2000.