

Medicinal Observations and Ethnobotanical Investigations of Bamboo Plants in Udupi Region

Mallya Suma¹, Lekshmy MS²

²Associate Professor, Department of Dravyaguna, Sri Dharmasthala Manjunatheshwara College of Ayurveda, Kuthpady, Udupi, 574118

¹PG Scholar, Department of Dravyaguna, Sri Dharmasthala Manjunatheshwara College of Ayurveda, Kuthpady, Udupi, 574118
sumamall002@gmail.com
+91984474002

Abstract: About: The floristic wealth of India is diverse and enormous. Many species of *Phyllanthaceae* used in medicine grow in all climatic conditions. Medicinal plant knowledge along with their therapeutic benefits, phenology, geological conditions is essential in current era. Materials and Methods: Field survey was conducted during flowering and fruiting season of *Phyllanthaceae* species in and around Udupi district plants identified, authenticated herbarium was prepared. Ethno - medico - botanical information about each species was collected by consulting local healers and detailed information listed by referring floras. Results: *P. emblica*, *P. acidus*, *P. reticulatus*, *P. urinaria*, *P. amarus*, *P. tenellus*, *P. debilis*, *P. rheedi*, *P. virgatus*, *P. lawii* and *P. myrtifolius* are major species found in study area. Discussion and Conclusion: Family *Phyllanthaceae*, having diverse medicinally beneficial plant species available wild, also found in cultivated status. 11 species of *Phyllanthaceae* have been documented from Udupi district, each having medicinal usage. Morphological, phytochemical, geological differences also found interesting among species.

Keywords: Ayurveda, *Phyllanthaceae*, ethno - medicinal uses, field survey

1. Introduction

Universal role of plants in the treatment of disease is exemplified by their employment in all major systems of medicine irrespective of the underlying philosophical premise¹. Plants sustain human life by providing the raw materials needed for living such as food, clothing, shelter, medicine, energy, along with rendering various ecosystem services to maintain the ecological balance². The floristic wealth of India is diverse and enormous. Western Ghats is one among biodiversity hotspots of India which is rich with diverse medicinal plants. In India it is estimated that there are around 17000 species of higher plants, among which 7500 are known for medicinal use³.

Phyllanthaceae considered as a subfamily of *Euphorbiaceae* many of which are possessing medicinal value, most numerous in tropical region⁴. Genus *Phyllanthus*, member of the flowering plant family *Phyllanthaceae*, includes more than 1000 species, distributed in tropical and subtropical regions⁵. Most of its species have *phyllanthoid* branching produce flowers only on the ultimate branchlets. Because of these branchlets often fall as a unit and resemble that of pinnate leaves, the flowers of the plant superficially appear to be borne on the leaves⁶. In some species of *Phyllanthus*, the ultimate branches are essentially leafless and flattened into strikingly leaf like cladodes; their homology with branches is clearly demonstrated by the cymules of flowers along the margins⁷. The difference between *Euphorbiaceae* and *Phyllanthaceae* are listed in table 1.

Table 1: Difference between *Euphorbiaceae* and *Phyllanthaceae*⁸

Plant characters	<i>Euphorbiaceae</i>	<i>Phyllanthaceae</i>
Ovary	Single ovule per locule	2 per locule
Style	3	2-5
Stigma	3, lobed (furtherlobed)	1
Fruit	Regma, capsule, drupe, or berry	Fruits usually capsules berries or drupes.
Seed	Arillate seeds	Non arillate seeds

Several *Phyllanthaceae* species are widely used in medicine, comprising diverse medicinal properties, and phytochemical constituents. Udupi is a coastal district of Karnataka, characterized by excessive humidity during greater part of the year. The purpose of this study is to conduct a comprehensive ethnobotanical survey of *Phyllanthaceae* species in the Udupi district, document their medicinal uses, and contribute to the knowledge of traditional medicine practices. This study is significant as it provides detailed ethno - medico - botanical information on *Phyllanthaceae* species, contributing to the preservation of traditional knowledge and potential discoveries in herbal medicine.

Hence with all this background a study has been planned to conduct a survey on medicinal plant species of *Phyllanthaceae* of Udupi district, along with their documentation on ethno - medico - botanical uses.

2. Materials and Methods

Field survey was conducted during flowering and fruiting season of *Phyllanthaceae* species in and around Udupi district plants identified, authenticated herbarium was prepared, referring flora and botanist opinion. Ethno - medico - botanical information about each species was

collected by consulting local healers and detailed information listed by referring floras, Materia medica as per standard guidelines^{9, 10}.

3. Results

Below listed plants have been found during systematic survey conducted in study area, which were documented along with their traditional medicinal benefits.

a) *Phyllanthusemblica* Linn.

A medium - sized deciduous tree with light grey bark found exfoliating in irregular patches. Leaves were subsessile, oblong - elliptic with flowers in axillary fascicles.^{11, 12} Drupaceous fruits and trigonous seeds were features. Planted in public parks, medicinal gardens. Fruits used in jaundice, dyspepsia, bacillary dysentery, eye trouble also as a gastrointestinal tonic. Juice with turmeric powder and honey is prescribed in diabetes insipidus. Seed as anti - asthmatic used in bronchitis. Leaf - juice is given in vomiting. Fruits are rejuvenator used in Indian system of medicine.^{13, 14} A decoction of powdered pericarp advised for peptic ulcer. Root bark as astringent, said to be beneficial in ulcerative stomatitis. Leaves recorded beneficial in conjunctivitis, inflammation, dyspepsia, diarrhoea and dysentery.

b) *Phyllanthusacidus* Skeels. / *Ciccaacida* (Linn.) Merrill

It is a small tree with ovate leaf and acute apex. Flowers reddish, minute with drupaceous, globose, 6 - 8 - angled fruit.^{15, 16} Usually cultivated in gardens for the edible fruits, which are used in pickling. It is used as liver tonic which improves appetite, useful in biliousness, constipation, vomiting, bronchitis.

c) *Phyllanthusreticulatus*. Poir^{17, 18}

It is an erect or straggling shrub having slender and drooping branchlets. Leaves found elliptic - orbicular, rounded at both ends, glabrescent; with 3 mm long petioles. Flowers located in axillary fascicles with globose, purple - black fruits. Commonly grown in hedges. *P. reticulatus* leaves are used in bleeding gums along with *Kankola* (*Piper cubeba*) and *Karpura* (*Cinnamon camphor*) as it stops the bleeding. The decoction of its root said to be beneficial in children for bronchitis. Stem decoction is also used in painful eyes and conjunctivitis. Fruits used in blood purification, also as a diuretic.

d) *Phyllanthusurinaria* L.

It is an erect, annual herb with sessile, closely placed, often overlapping leaves. Male flowers pedicellate whereas female flowers axillary, solitary. Fruits were capsule with transversely ridged seeds¹⁹. Found common in wastelands and as weed in gardens. It is an excellent diuretic whereas leaves are eaten by cattle. The juice of the leaves administered with coconut milk as an appetizer to children.

e) *Phyllanthustenellus* Roxb²¹.

It is an annual glabrous herb, with elliptic to obovate, acute leaf. Flowers found in axillary fascicles having filiform Capsule and trigonous seeds. Native to Mascarene Islands, but now well naturalized, found everywhere. It is used to treat inflammation and liver diseases.

f) *Phyllanthusamarus* Schumach & Thonn.

It is an erect herb with subsessile, closely placed and often overlapping, oblong to obovate leaves. Flowers found in fascicles with depressed globose capsule²². Found as a weed in cultivated and waste lands. Used as a single drug in the treatment of jaundice as decoction. Used in prescriptions for dyspepsia, indigestion, chronic dysentery, urinary tract diseases, diabetes, skin eruptions. It is also useful in jaundice, diarrhoea, dysentery, intermittent fevers, diseases of the urino - genital system, scabies, ulcers.

g) *Phyllanthusmyrtifolius* (Wight) Muell. Arg.

It is a rigid shrub with angled branches and narrowly oblanceolate leaf. Flowers pinkish or greenish, found in bracteate axillary fascicles. Fruits are filiform capsules having trigonous seeds. Native to Sri Lanka; often cultivated in garden as a hedge plant²³.

h) *Phyllanthus debilis* Klein ex Willd.

It is an erect herb with closely placed, elliptic - acute leaves and short petiole. It has very short, flowers in fascicles. Found as a common weed in paddy fields after harvest.²⁴

The herb said to be bitter in taste, and is reported to possess astringent, diuretic, febrifugal and antiseptic properties. It is used in stomach troubles such as dyspepsia, colic, diarrhoea and dysentery. It is also employed in swelling and diseases of urinogenital system²⁵.

i) *Phyllanthusrheedi* Wight

It is a slender, erect glabrous herb with branched stem. Leaves were elliptic or ovate and flowers found in axillary fascicles with oblate capsule having trigonous, 6 - 7 - ridged seed inside. Found in grass lands and commonly used in dysentery²⁶

j) *Phyllanthusvirgatus* G. Forst

It is an erect herb or undershrub with woody rootstock and long flattened branches. Leaves were subsessile, closely placed and often overlapping. Male flowers few, minute, subsessile whereas female on filiform pedicels. Depressed - globose, long - stalked capsule and red tubercular seeds are characteristic features. Commonly found in waste lands, hedges.

Plant said to be antiseptic, wound healing. Fresh leaves, bruised in butter milk, are used as a wash for itching. Fresh leaves, flower and fruits along with cumin seeds and sugar, are used in gonorrhoea. Root application said to be beneficial in mammary abscesses. The juice of the leaves is employed in eye diseases²⁷.

k) *Phyllanthuslawii* J. Graham

Thorny bushy shrub with purplish branches armed with stipular tubercles. Leaves long, elliptic - oblong, shortly stalked. Flowers in axillary fascicles with globose capsule. Frequently found on the banks and in the beds of rocky rivers. Its branches are used for making baskets²⁸.


Thus, totally around 11 species have been found in Udupi district, among which herbarium and live photographs of 8 species have been documented.




Figure 1: Plant Species belonging to Family *Phyllanthaceae*




2/a. *Phyllanthusemblica* Linn.



22/b. *Phyllanthus tenellus* Roxb



2/c. *Phyllanthusurinaria* Linn.



2/d. *Phyllanthusreticulatus*. Poir




Figure 2: Herbarium of Plant Species belonging to Family Phyllanthaceae

4. Discussion

India treasure house of medicinal plant species, has diverse sources which are used in various therapeutics²⁹. *Phyllanthaceae* subfamily of *Euphorbiaceae* found numerous in tropical region. Genus *Phyllanthus* a member of the flowering plant family *Phyllanthaceae*, includes more than 1000 species showing characteristic *phyllanthoid* branching producing flowers only on the ultimate branchlets.

Ethno medico botanical survey has its application in many fields such as biodiversity, conservation strategy, human health, sustainable harvesting procedures etc³⁰.

Survey revealed that many plants of *phyllanthaceae* species have ethno - medicinal uses. Among these *Phyllanthus emblica* Linn. a medium sized tree, fruits of which are most widely used in Ayurveda for rejuvenation. *Phyllanthus acidus* Skeels. is a small tree having edible fruits, used in culinary preparations

Phyllanthusreticulatus Poir. an erect or straggling shrub aerial part of which are used in wound healing. *Phyllanthusurinaria* an annual herb known as an excellent diuretic, whereas *Phyllanthus amarus* Schumach & Thonn, and *Phyllanthus debilis* effective medicine used in the treatment of liver diseases. *Phyllanthus rheedi* Wight is given in dysentery and *Phyllanthus virgatus* G. Forst used as antiseptic in gonorrhea. Others like *Phyllanthus lawii* J. Graham and *Phyllanthus myrtifolius* (Wight) Muell. Arg. have commercial importance³¹.

11 species of *Phyllanthaceae* have been documented from Udupi district, each having medicinal usage. Morphological, phytochemical, geological differences also found interesting among species.

5. Conclusion

Family *Phyllanthaceae*, having varied medicinally beneficial plant species available wild, also found in cultivated status. Morphological characters, medicinal benefits enlisted above contributory in further new drug research. This study not only documents the medicinal properties of *Phyllanthaceae* species in the Udupi district but also underscores the importance of preserving traditional knowledge for future pharmacological explorations.

References

- [1] Sharma Rakesh K, Arora Rajesh. Herbal drugs a twenty first century perspective. Jaypee Brothers Medical Publishers (P) Ltd.2006.9p
- [2] Swagata Tavhare D, Nishteshwar K. Collection practices of medicinal plants - Vedic, Ayurvedic and Modern perspectives. International journal of Pharmaceutical and biological Archives.2014; 5 (5); 54 - 61
- [3] Divya et al. IUCN Red listed medicinal plants of Siddha. J Res Sid Med.2018; 1 (1); 15 - 22
- [4] Nadakarni KM, Indian Materia Medica; Popular Prakashan; Mumbai 2010; 948
- [5] Patel JR, Tripathi P, Sharma V, Chauhan NS, Dixit VK. *Phyllanthus amarus*: ethnomedicinal uses, phytochemistry and pharmacology: a review. J Ethnopharmacol.2011; 138 (2); 286–313.
- [6] Madamanchi Geethanjali, Shih - Torng Ding. A review of the Phytochemistry and Pharmacology of *Phyllanthus urinaria*L. Front Pharmacol.2018; 1109
- [7] Aparupa Bose, Mazumdar Ghosh, Anindita Banerjee, Sharmila Chattopadhyaya. An insight into the potent medicinal plant *Phyllanthus amarus*Schum. And Thonn. The Nucleus.2022; 437 - 72
- [8] Adel Aziz Fayed, Mohammed Soliman, Ahamed Faied, Mona Hassan. Taxonomic evaluation of *Euphorbiaceaesensulato* with special reference ti*Phyllanthaceae* as a new family to the flora of Egypt. Biological Forum - An International Journal.2019; 11 (1); 47 - 64
- [9] Chinmay Rath, Bonthu Susmitha, Ashish Tripathi K, Mukhesh Chincholkar B, Anupam Mangal K, Narayanam Srikanth. Medico ethno botanical Survey; An overview of CCRAS contributions. JDRAS.2017; 2 (3); 188 - 240
- [10] Srikanth N, Singh S, Sharma BS, Khanduri S, Singh R, Maheshwar T. Tribal health care research program: an overviewof CCRAS contributions. I Drug Res Ayurvedic sciences.2017; 2 (2): 118 - 48
- [11] Warriar PK. Indian Medicinal Plants. Orient Longman Private Ltd.1994.412p
- [12] Bhat Gopalakrishna K. Flora of Udupi. Indian Naturalist.2003.576p
- [13] Hegde PL, Harini A. A Textbook of DravyagunaVijnana. Chaukhamba Publications.2021.383p
- [14] Kallianpura Supriya, Gokarna Rohit A. A text book of DravyagunaVijnana. Chaukhamba Orientalia.2019, 198p
- [15] Tan SP, Tan ENY, Lim QY, Nafiah MA. *Phyllanthus acidus* (L.) Skeels: A review of its traditional uses, phytochemistry, and pharmacological properties. J Ethnopharmacol.2020; 253 - 112610.
- [16] Kirtikar KR, Basu B. Indian medicinal plants. International Book distributors.2006.1369p
- [17] Kharat A. Review on *Kirganelia reticulata*. Journal of Drug Delivery and Therapeutics.2013; 3 (4); 208–12.
- [18] Vaidya Bapalal G. Nighantu Adarsha. Chaukhamba Bharati Academy.1998.667p
- [19] Madamanchi Geethanjali, Shih Torng Ding. A review of the Phytochemistry and Pharmacology of *Phyllanthus urinaria* L. Front Pharmacol.2018; 9 - 1109
- [20] Khare CP. Indian medicinal Plants an illustrated dictionary. Springer.2007.297p
- [21] Tassia Campos de Lima e Silva, Jorge Veras Filho, Elba Lucia Cavalcanti de Amorim, Ivone Antonia de Souza, UlyssesPaulino de Albuquerque, EdnaldoCavalcante de Araujo. Acute toxicity study of stone breaker (*Phyllanthus tenellus*Roxb.). Rev Cienc Farm Basica Apl.2012; 33 (2); 205 - 210
- [22] Bose Mazumdar Ghosh A, Banerjee A, Chattopadhyay S. An insight into the potent medicinal plant *Phyllanthus amarus* Schum and Thonn. Nucleus (Calcutta).2022; 65 (3); 437–72.
- [23] Chang CW, Lin MT, Lee SS, Liu KC, Hsu FL, Lin JY. Differential inhibition of reverse transcriptase and cellular DNA polymerase - alpha activities by lignans isolated from Chinese herbs, *Phyllanthus myrtifolius* Moon, and tannins from *Lonicera japonica* Thunb and *Castanopsishystrix*. Antiviral Res.1995; 27 (4); 367–74.
- [24] Perera D, Soysa P, Wijeratne S. Polyphenols contribute to the antioxidant and antiproliferative activity of *Phyllanthus debilis* plant in - vitro. BMC Complementary and Alternative Medicine.2016; 16 (1); 339.
- [25] Wanniarachchi K, Peiris D, Ratnasooriya W. Antihyperglycemic and hypoglycemic activities of *Phyllanthus debilis* aqueous plant extract in mice. Pharmaceutical Biology.2009, 1 (47); 260–65.
- [26] Suresh V, Asha VV. Preventive effect of ethanol extract of *Phyllanthus rheedii* Wight. on D - galactosamine induced hepatic damage in Wistar rats. J Ethnopharmacol.2008; 116 (3); 447–53.
- [27] Arshya Hashim, M Salman Khan, Mohd Sajid Khan, Mahd Hassan Baig, Saheem Ahmad. Antioxidant and α - Amylase Inhibitory Property of *Phyllanthus*

virgatus L: An In Vitro and Molecular Interaction Study. BioMed Research International.2013; 729393

- [28] MukharjeePulok K. Quality Control of Herbal Drugs. Business Horizons.2002.68p
- [29] Manju, Tasavur Ahad. Ethnobotanical studies - Importance and conservation strategies. Biological Diversity Current status and Conservation Policies.2021, 24 - 36
- [30] Chandrashekar KS, Saha S, Subraya PK. Analgesic Activity of Phyllanthus lawii Extract in Swiss Albino Mice. Pharmaceutical Crops.2011, 2: 8 - 1

Author Profile

- [1] Dr Suma Mallya MD, PhD (Dravyaguna) Associate professor, Department of Dravyaguna; Sri DharmasthalaManjunatheshwara college of Ayurveda Kuthpady, Udupi, Karnataka. Mobile: 9844740027; E mail: sumamallya[at]gmail.com
- [2] Dr Lekshmy MS, PG Scholar, Department of Dravyaguna Sri DharmasthalaManjunatheshwara college of Ayurveda, Kuthpady, Udupi, Karnataka; Mobile: 9496611844; E mail: lekshmyana[at]gmail.com